

P25. Shock waves induce angiogenesis via exosome release

Graber M¹, Pölzl L¹, Lobenwein D¹, Graber M¹, Blumer M¹, Kirchmair E¹, Kirchmair R¹, Paulus P¹, Davidson S¹, Grimm M¹, Tepeköylü C¹, Holfeld J.¹

Email: graber.michael@gmx.at

1. Medical University of Innsbruck, Department of Cardiac Surgery

Introduction	The mechanism of mechanotransduction of shock wave therapy (SWT) remains unknown. We hypothesized that SWT induces exosome release and thus exerts its angiogenic effects.
Material and Method	Human umbilical vein endothelial cells (HUVECs) were treated with SWT. Exosomes were isolated from the supernatant and analyzed by nanoparticle tracking analysis. The angiogenic potential was analyzed in vitro and exosomes were injected into subcutaneously implanted matrigel plugs in nude mice. Perfusion of the plugs was measured via Laser Doppler perfusion imaging (LDPI) and quantified histologically.
Results	Supernatants of treated HUVECs showed significantly higher concentrations of exosomes. Released exosomes enhanced tube formation and endothelial cell proliferation. Exosome-release inhibitor GW4869 abolished the angiogenic effects of SWT. Injection of exosomes into subcutaneously implanted matrigel plugs resulted in higher perfusion and increased number of capillaries and arterioles.
Discussion	We show for the first time how the mechanical stimulus of SWT is translated into a biological response. SWT causes exosome release which have and highly angiogenic potential in vitro and in vivo.
Conclusion	Exosomes released by SWT might develop a therapeutic treatment option for ischemic heart disease.